

Candida Antibodies & Immune Complexes



**Genova
Diagnostics®**

Innovative Testing for Optimal Health

63 Zillicoa Street
Asheville, NC 28801
© Genova Diagnostics

Patient: **SAMPLE
REPORT**

Age: 74
Sex: M
MRN:

Order Number:

Completed: November 19, 2007
Received: November 17, 2007
Collected: November 14, 2007

Anti-Candida Antibody

Reference Range

Candida Immune Complex (serum)	<input type="text"/>	<input type="text" value="1.2"/>	<= 0.9
Candida Antibody, IgA (serum)	<input type="text"/>	<input type="text" value="22"/>	<= 10
Candida Antibody, IgG (serum)	<input type="text"/>	<input type="text" value="11"/>	<= 10
Candida Antibody, IgM (serum)	<input type="text" value="4"/>	<input type="text"/>	<= 10

Commentary

The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. Unless otherwise noted with * as cleared by the U.S. Food and Drug Administration, assays are For Research Use Only.

Commentary is provided to the practitioner for educational purposes, and should not be interpreted as diagnostic or treatment recommendations. Diagnosis and treatment decisions are the responsibility of the practitioner.

Candida immune complexes contain IgG Candida antibodies, Candida antigen and fragments of complement. Elevations of these complexes correlate with the clinical symptoms related to chronic Candidiasis, and diminish with effective Candida treatment.

Elevated levels of IgA Candida antibody titers are indicative of local mucosal infections, including the vagina, skin, urethra, and GI tract.

Chronic candidiasis can be positively diagnosed by the determination of high serum levels of IgG and IgM antibodies against *Candida albicans*. High levels of IgG antibody titers can be indicative of past or ongoing Candida infection.